

APPENDIX D. PRODUCT LABELS

<p>Biological Insecticide</p> <p>Foray® 76B</p> <p>Flowable Concentrate</p> <p>For the control of Lepidopterous Larvae</p> <p>ACTIVE INGREDIENT: <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i>, strain ABTS-351, fermentation solids, spores, and insecticidal toxins 18.44% Other Ingredients: 81.56% Total: 100.0%</p> <p>Potency: 16,700 Cabbage Looper Units (CLU)/mg of product (equivalent to 76 billion CLU/GAL.). The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.</p> <p>EPA Reg. No. 73049-49 EPA Est. No. 33762-IA-001</p> <p style="text-align: right;">LIST NO. 60176</p> <hr/> <p>INDEX:</p> <ul style="list-style-type: none"> 1.0 First Aid 2.0 Precautionary Statements <ul style="list-style-type: none"> 2.1 Hazards to Humans and Domestic Animals 2.2 Personal Protective Equipment (PPE) 2.3 Agricultural Use Requirements 2.4 Non-Agricultural Use Requirements 2.5 User Safety Recommendations 2.6 Environmental Hazards 3.0 Directions for Use 4.0 Storage and Disposal 5.0 Agricultural Use Requirements 6.0 Non-Agricultural Use Requirements 7.0 Application 8.0 Mixing 9.0 Spray Volumes 10.0 General Agricultural Use Instructions 11.0 Table 1 12.0 General Non-Agricultural Use Instructions <ul style="list-style-type: none"> 12.1 Ground Application 12.2 Aerial Application 13.0 Table 2 14.0 Notice of Warranty <hr/> <p>KEEP OUT OF REACH OF CHILDREN CAUTION</p>	<p>1.0</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="2" style="text-align: center;">FIRST AID</th> </tr> </thead> <tbody> <tr> <td style="width: 50px; vertical-align: top; padding: 5px;">If on skin or clothing</td> <td style="padding: 5px;"> <ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. </td> </tr> <tr> <td style="width: 50px; vertical-align: top; padding: 5px;">If in eyes</td> <td style="padding: 5px;"> <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. </td> </tr> <tr> <td colspan="2" style="text-align: center;">HOT LINE NUMBER</td> </tr> <tr> <td colspan="2" style="padding: 5px;"> <p>Have the product container with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-315-9819 for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-323-9597.</p> </td> </tr> </tbody> </table> <hr/> <p>2.0 PRECAUTIONARY STATEMENTS</p> <p>2.1 HAZARDS TO HUMANS AND DOMESTIC ANIMALS</p> <p>CAUTION</p> <p>Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.</p> <p>2.2 Personal Protective Equipment (PPE)</p> <p>Applicators and other handlers must wear:</p> <ul style="list-style-type: none"> Long-sleeved shirt Long pants Waterproof gloves Shoes plus socks <p>Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.</p> <p>2.3 Agricultural Use Requirements</p> <p>Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic reactions. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.</p> <p>IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must provide all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.</p> <p>2.4 Non-Agricultural Use Requirements</p> <p>Mixer/loaders and applicators not in enclosed cabs or aircraft must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.</p>	FIRST AID		If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 	If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 	HOT LINE NUMBER		<p>Have the product container with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-315-9819 for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-323-9597.</p>	
FIRST AID											
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 										
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 										
HOT LINE NUMBER											
<p>Have the product container with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-315-9819 for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-323-9597.</p>											

CONTINUED

2.5 User Safety Recommendations Users should: <ul style="list-style-type: none">• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.• Remove clothing immediately if pesticide gets inside. Wash thoroughly and put on clean clothing.• Remove PPE immediately after handling the product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.	5.0 AGRICULTURAL USE REQUIREMENTS Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: <ul style="list-style-type: none">• Coveralls• Waterproof gloves• Shoes plus socks
2.6 Environmental Hazards For terrestrial agricultural uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. This product must not be applied aerially within 1/4 mile of any habitats of threatened or endangered lepidoptera. No manual application can be made within 300 feet of any threatened or endangered lepidoptera.	
3.0 DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.	
4.0 STORAGE AND DISPOSAL Do not contaminate water, food or feed by storage or disposal of waste. Pesticide Storage: Store in a cool, dry place. Keep containers tightly closed when not in use. Store in temperatures above freezing and below 25° C (77° F). Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry). Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Once cleaned, offer container for recycling, if available. If recycling is not available, puncture and dispose of container in a sanitary landfill or by other procedures approved by state and local authorities.	6.0 NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. For ground applications only. Exposure of unprotected persons can be mitigated by direct spraying. Spray should be allowed to dry undisturbed.

- 7.0 **APPLICATION**
Apply Foray 76B by ground or aerial equipment undiluted or with quantities of water sufficient to provide thorough coverage of plant parts to be protected. The amount of water needed per acre will depend upon crop size, weather, spray equipment, and local experience.
Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower/treatment coordinator are responsible for considering all of these factors when making decisions.
- 8.0 **MIXING**
Shake or stir Foray 76B before use. Fill spray or mixing tank half full of water. Begin agitation and pour Foray 76B into water while maintaining continuous agitation. Add other spray material (if any) and balance of water. Agitate as necessary to maintain suspension. Do not allow diluted mixture to remain in the tank for more than 72 hours.

To improve weather-fastness of the spray deposits for hard to wet crops, such as cole crops, use a spreader-sticker approved for use on growing crops. Combinations with commonly used spray tank adjuvants are generally not deleterious to Foray 76B, if the mix is used promptly. Before mixing in the spray tank, identify possible problems with physical compatibility by mixing all components in a small container in proportionate quantities.

9.0 SPRAY VOLUMES

Ground Application: Use amount of Foray 76B, as indicated in the tables that follow, in ground equipment with quantities of water sufficient to provide thorough coverage of plant parts to be protected. The amount of water needed per acre will depend upon crop size, weather conditions, spray equipment used and local experience.

Aerial Application: Use amount of Foray 76B, as indicated in the tables that follow, in aerial equipment undiluted or with quantities of water sufficient to provide thorough coverage of plant parts to be protected. In the western U.S., use a normal minimum of 5-10 gallons per acre; in the eastern regions, use a normal minimum of 2-3 gallons per acre. The minimum amount of water needed per acre will depend upon crop size, weather conditions, spray equipment used and local experience.

10.0 GENERAL AGRICULTURAL USE INSTRUCTIONS

Foray 76B is a biological insecticide for the control of lepidopterous larvae. It contains the spores and endotoxin crystals of *Bacillus thuringiensis kurstaki*. Foray 76B must be ingested by the larvae to be effective. For consistent control, apply at first sign of newly hatched larvae (1st and 2nd instar larvae). Susceptible larvae that ingest Foray 76B cease feeding within a few hours and die within 2-5 days.

Foray 76B may be applied up to and on the day of harvest. For maximum effectiveness, follow the instructions listed below:

Monitor fields to detect early infestations.

Apply Foray 76B when eggs start hatching and larvae are small (early instars) and before significant crop damage occurs. Larvae must be actively feeding to be affected.

Repeat applications every 3 to 14 days to maintain control and protect new plant growth. Factors affecting spray interval include rate of plant growth, weather conditions, and reinfections. Monitor populations of pests and beneficials to determine proper timing of applications.

Under conditions of heavy pest pressures or when large worms are present use the higher rate, shorten the application interval, and/or improve spray coverage to enhance control. When these conditions are present, consider use of a contact insecticide to enhance control. Thorough coverage is essential for optimum performance. Ground applicators equipped with directed drop nozzles can improve coverage.

11.0 Table 1.

Crop	Pests	Rate ¹ (fl. oz./acre)
Forests, Shade Trees, Ornamentals, Shrubs, Sugar Maple Trees, Ornamental Fruit, Nut & Citrus Trees ²	Gypsy Moth ² Elm Spanworm Spruce Budworm Browntail Moth Douglas Fir Tussock Moth Cone worm Buck Moth	13.5 - 67.5
	Tussock Moth Pine Butterfly Bagworm Leafroller Tortrix Mimosa Webworm Tent Caterpillar Jackpine Budworm Blackheaded Budworm Saddled Prominent Saddleback Caterpillar Eastern & Western Hemlock Looper Orangetripped Oakworm Satin Moth	10.0 - 27.0
	Redhumped Caterpillar Spring & Fall Cankerworm California Oakworm Fall Webworm	7.0 - 13.5

Special Instructions

¹Use the higher recommended rates on advanced larval stages or under high density larval populations.

²In treating gypsy moth infested trees and shrubs in urban, rural, and semi-rural areas, exposure of non-target vegetation including, but not limited to, native and ornamental species and food or feed crops is permitted.

This product can be mixed and used with other pesticides only in accordance with the most restrictive of label limitations and precautions. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rates may be exceeded.

12.0 GENERAL NON-AGRICULTURAL USE INSTRUCTIONS

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on trees being grown for sale or other commercial use, or for commercial seed production, or for the production of timber or wood products, or for research purposes except for wide-area public pest control programs sponsored by government entities, such as mosquito abatement, gypsy moth control, and Mediterranean fruit fly eradication.

Do not apply this product through any type of irrigation system.

Foray 76B contains the spores and endotoxin crystals of *Bacillus thuringiensis kurstaki*. Foray 76B is a stomach poison and is effective against lepidopterous larvae. After ingestion, larvae stop feeding within hours and die 2-5 days later. Maximum activity is exhibited against early instar larvae. Apply Foray 76B by ground or aerial equipment.

CONTINUED

Shake or stir Foray 76B before use. Add some water to the mix tank, pour the specified amount of Foray 76B into the tank, and then add the remaining amount of water to obtain the proper mix ratio. Agitate as necessary to maintain the suspension. Do not allow diluted mixture to remain in the tank for more than 72 hours.

12.1 Ground Application

Use an adequate amount of tank mix to obtain thorough coverage without excessive run off. Use the indicated recommended per acre rates of Foray 76B in up to the following amounts of water:

High volume hydraulic sprayers	100 gallons
Mist blowers	10 gallons

12.2 Aerial Application

Apply Foray 76B, either alone or diluted with water, aerially at the rates shown in the application rates table. Spray volumes of 28-67.5 fluid ounces of product per acre give optimum coverage.

13.0 Table 2.

Crop	Pests	Rate ¹ (fl. oz./acre)
Forests, Shade Trees, Ornamentals, Shrubs, Sugar Maple Trees, Ornamental Fruit Nut & Citrus Trees ²	Gypsy Moth ² Elm Spanworm Spruce Budworm Browntail Moth Douglas Fir Tussock Moth Coneworm Buck Moth Tussock Moth Pine Butterfly Bagworm Leafroller Tortrix Mimosa Webworm Tent Caterpillar Jackpine Budworm Blackheaded Budworm Saddled Prominent Saddleback Caterpillar Eastern & Western Hemlock Looper Orangetripped Oakworm Satin Moth Redhumped Caterpillar Spring & Fall Cankerworm California Oakworm Fall Webworm	13.5 - 67.5 13.5 - 50.5 10.0 - 27.0 7.0 - 13.5

Special Instructions

¹Use the higher recommended rates on advanced larval stages or under high density larval populations.

²In treating gypsy moth infested trees and shrubs in urban, rural, and semi-rural areas, exposure of non-target vegetation including, but not limited to, native and ornamental species and food or feed crops is permitted.

14.0 NOTICE OF WARRANTY

To the extent consistent with applicable law, seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risk of use, storage or handling not in strict accordance with accompanying directions.

Foray is a registered trademark of Valent BioSciences Corporation.



870 TECHNOLOGY WAY
LIBERTYVILLE, IL 60048 USA
PH: 800-323-9597

04-6159/R5 ©Valent BioSciences Corporation, August 2009

HERCON®
DISRUPT® II
GYPSY MOTH MATING DISRUPTANT

Population Suppressant

HERCON® DISRUPT® II Gypsy Moth is a controlled-release pheromone formulation designed to lower incidence of gypsy moth, *Lymnia xantharia*, mating by disrupting normal male flight orientation to females. This reduction in mating will help suppress the larval caterpillar population that causes damage by feeding on the leaves of hardwoods and evergreens.

ACTIVE INGREDIENTS:

(Z)-7,8-epoxy-2-methyloctadecane	17.9 %*
OTHER INGREDIENTS	82.1 %
TOTAL	100.0 %

CONTENTS:

MINIMUM NET WEIGHT:	[KG] [lb]*
* 8.5 kg [18.7 lb] of product will treat 50 acres at 30.4 g/A./acre	

KEEP OUT OF REACH OF CHILDREN
C A U T I O N

Read Directions and Precautionary Statements Before Use

FIRST AID:

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

IF IN EYES:

- Hold eye open and gently wash with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor immediately for treatment advice.

IF ON SKIN:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor immediately for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor immediately for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the National Pesticide Information Center (NPIC) at 1-800-852-3738 seven days a week, 8:30 am to 4:30 pm Pacific Time (NPIC Web site: www.npic.orst.edu). After 4:30 pm call your poison control center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes and mouth. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco products or using the toilet. Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

ENVIRONMENTAL HAZARDS: For terrestrial uses: Do not apply directly to water or to areas where surface water is present nor to intertidal areas below the mean high water mark, except under forest canopy. Do not contaminate water when disposing of equipment washwaters or rinseate.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply this product up to two weeks before adult gypsy moth emergence. Depending on the gypsy moth population densities apply 30 gm (170 gm (6 oz) of product), 15 gm (85 gm (3 oz) of product) or 6 gm (34 gm (1.2 oz) of product) of active ingredient per application per acre. Apply 15 gm and 6 gm of active ingredient in low density gypsy moth populations. Consult your state or local authorities for determining gypsy moth population levels in your area. To ensure proper rate and method of application, make application by or under the supervision of qualified a person.

Apply a second application if adult gypsy moth emergence is extended or delayed, otherwise one application lasts the entire season. Use an inert sticker material with DISRUPT® II to hold flakes on treated foliage or plant parts. The Hercon applicator is specifically designed to mix the proper amount of DISRUPT® II flakes and inert sticker at the time of application. Use in areas such as forest, residential, municipal and shade tree areas, recreational areas such as campgrounds, golf courses, parks and parkways, ornamental, shade tree plantings, shelter belts and rights of way and other easements.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in sealed containers in a cool dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Plastic Bails: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of bag in a sanitary landfill or by incineration if, if allowed by state or local authorities, by burning. If burned stay out of smoke. Metal Containers (outside containers): Non-refillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

WARRANTY AND DISCLAIMER STATEMENT
To the fullest extent permitted by law, Hercon Environmental warrants that this material conforms to the chemical description on the label. Manufacturer neither makes nor authorizes any agent or representative to make any other warranty of fitness or on merchantability, guarantee or representation, expressed or implied concerning this material. Manufacturer's maximum liability for breach of this warranty shall not exceed the purchase price of this product. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this material not in conformance with the label.
Aberdeen Road Company dba HERCON ENVIRONMENTAL
P.O. Box 335 Emigsville, PA 17318-0335 EPA Est. No. 8780-PA-01
P.O. Box No. 8780-PA-01 Questions? Call 1-866-4-HFCON
© HERCON, BIO-FLAKE and DISRUPT are registered trademarks of Aberdeen Road Company. REV 06/10/09

ACTIVE INGREDIENT:	Disparlure, (Z)-7,8-Epoxy-2-methyloctadecane.....	13.00%
OTHER INGREDIENTS.....		87.00%
TOTAL.....		100.00 %

SPLAT

GMTM

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID STATEMENT

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Wash skin immediately with soap and water, then rinse with plenty of water for 15-20 minutes.

IF SWALLOWED:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

National Pesticides Information Center: Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: wwwnpic.orst.edu).

See side/back panel for additional precautionary statements

Net Contents: _____	Lbs _____	Kg _____
Batch Number: _____		

EPA Reg. No. 80286-4
EPA Est. No. 80286-CA-004



HAZARDS TO HUMANS AND DOMESTIC ANIMALS, CAUTION: Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before use.		with paint mixing paddle or similar device works best. Avoid mechanical mixing with sharp blades which come into contact with the plastic container as this can introduce sharpness to the SPLAT GM formulation, which might interfere with product flowability and application.			
PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.		SPLAT GM is applied either aerially or by ground with conventional application systems pressurized either by centrifugal pump or positive displacement pump; pressurized gas cylinders, or a combination of both. The flow of SPLAT GM product can be controlled with agricultural-type solenoids. Calibrate applications based on volume output and application speed and water treated. Do not apply this product through any type of irrigation system. SPLAT GM can also be applied manually using other devices like a metered dosing gun. To ensure proper application, consult your pest control advisor, distributor, or ISCA Technologies, Inc.			
ENVIRONMENTAL HAZARDS: For terrestrial uses: Do not apply directly to water, or areas where surface water is present or to potential areas below the mean high water mark. Incidental applications to water under the forest canopy are not considered to be a direct application to water, but should be avoided if practicable. Spray valves must be shut off when passing over surface water or under the canopy. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.		For ground applications, SPLAT GM droplet size will vary depending on pest population and duration of control desired. A dense infestation will require a higher density of point sources to achieve optimal mating disruption. With aerial applications, there is less control over the variation in droplet size and there will be a range from approximately 300 to 2000 microns in size. When SPLAT GM is sprayed on the tree canopy at 46 grams per acre (6 grams Al per acre), the range in droplet size will usually be 300 to 1000 microns. This will provide area-wide, season-long management of low-density populations. To ensure proper application, please consult your pest control advisor, distributor, or ISCA Technologies, Inc.			
DIRECTIONS FOR USE		STORAGE AND DISPOSAL			
<p>GENERAL SPLAT GM uses the pheromone of the gypsy moth (<i>Lymantria dispar</i>), to disrupt adult moth mating and thus reduce larval damage to trees.</p> <p>TARGET PEST Gypsy Moth(<i>Lymantria dispar</i>)</p> <p>AREA-WIDE CONTROL This product is only for use in wide-area gypsy moth control programs sponsored by governmental entities. Areas of application include, but are not limited to: forests, residential, municipal, and shade tree areas, recreational areas such as campgrounds, golf courses, parks, and parkways; ornamental and roadside trees; shelterbelts, rights-of-way, and other easements. A near-wide or wide-area application of SPLAT GM is most effective. The efficacy of any mating disruption strategy will be reduced by the influx of adults from surrounding areas. It is for this reason that mating disruption works increasingly well with larger acreages, where the ratio of exposed borders to overall area treated is reduced. Always make application of SPLAT GM in consultation with your Pest control advisor, distributor, or ISCA Technologies, Inc.</p> <p>No intentional food uses are permitted. However, an exemption from the requirement of a tolerance is established for residues of (Z)-7-heptenoic-2-methyldecane (Disparlure) on all food and feed crops that occur when it is used to treat trees, shrubs, and pastures and such use results in unintentional spray and drift to non-target vegetation including non-food, food, and feed crops.</p> <p>MONITORING Monitor for pests on a regular basis so that timely intervention with insecticides or alternative controls are possible. Use gypsy moth pheromone-baited monitoring traps made by ISCA Technologies or other suitable traps to monitor the presence of male moths, their emergence and their movement. Place monitoring traps before applying SPLAT GM to determine the infestation level. Continue monitoring throughout the season to evaluate efficacy and longevity of the pheromone application. Monitor treated plots for the target pest. Implement supplemental control measures if the pest population densities are above economic threshold levels.</p> <p>APPLICATION TIMING Apply SPLAT GM anytime in the two weeks preceding male flight and adult activity. Additional applications may be made if populations exceed the economic threshold. Consult your pest control advisor, distributor, or ISCA Technologies, Inc for proper application timing.</p> <p>APPLICATION CONDITIONS Apply SPLAT GM when the ambient air temperature is above 50°F. SPLAT GM will cure within 3-4 hours following application, after which it will become rain fast and UV resistant. Do not apply if rain is expected within 1-2 hours of application or the temperature is outside of this range.</p> <p>APPLICATION RATE The application rate of SPLAT GM ranges from 22 grams to 230 grams of undiluted product per acre. This is equivalent to 3 grams to 30 grams of pheromone active ingredient (Al) per acre. Do not exceed 150 grams of Al per acre per year for multiple applications.</p> <p>PREPARATION FOR USE Always check SPLAT GM 24 hours before using. If SPLAT GM has been stored in a refrigerated area and has hardened, allow at least 48 hours at room temperature to achieve a workable consistency. Always mix SPLAT GM thoroughly before use; a drill</p>		<p>PESTICIDE STORAGE Do not contaminate water, food or feed by storage or disposal. Store product sealed and refrigerated, if possible. If refrigerated, do not store with food. Do not exceed 75°F for long-term storage. Avoid freezing. In case of leak or spill, wipe with paper towel and dispose of waste when product hardens.</p> <p>PESTICIDE DISPOSAL Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.</p> <p>CONTAINER DISPOSAL Nonrefillable container. Do not refill or reuse this container. Triple rinse (or equivalent) promptly after emptying. Then offer for recycling if available, or puncture and dispose of in sanitary landfill, or by incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.</p>			
<p>WARRANTY AND LIMITATION OF DAMAGES</p> <p>Read all directions carefully. All statements concerning the use of this product apply only when used as directed. ISCA Technologies warrants that this material conforms to the specifications on the label and is reasonably fit for the intended purpose referred to on the label. To the extent consistent with applicable law, ISCA Technologies makes no other express or implied warranty of merchantability or fitness for a particular purpose or any other express or implied warranty.</p>					
<p>Manufactured by: ISCA Technologies, Inc. 1201 Spring St. Riverside, CA 92507 (951) 686-5008 info@isca-tech.com</p> <p>ISCA TECHNOLOGIES</p>					